## McHUGH'S LANDSCAPING

PO BOX 908 DELRAN, NJ 08075 (856) 461-5154 FAX- (856) 461-6135

DATE: 3/25/19

### **PESTICIDE NOTIFICATION -**

CUSTOMER: DELRAN BOE

ADDRESS: 52 HARTFORD RD., DELRAN, NJ 08075

TREATED AREAS: ATHLETIC FIELDS AT DHS, DMS & DIS

DATE & TIME OF APPLICATION: FRIDAY, 3/29/19 7:00 AM TO 11:00 AM FIELDS MUST BE CLOSED UNTIL 2:00 PM

RE-ENTRY DATE & TIME: FRIDAY, 3/29/19, 2:00 PM

MATERIALS USED: 'DIMENSION PLUS FERTILIZER'
EPA# 10404-86

JAMES J. MCHUGH, (609) 352-5755 MCHUGH'S LANDSCAPING

PLEASE RETAIN FOR YOUR RECORDS.

# **DIMENSION\* 0.15% Plus Fertilizer**

NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. Cool-Season Grasses

In New York State this product may only be used by commercial applicators and at no more than 333 lb (0.5 lb of active ingredient) per acre per year (or 7.6 lbs product per 1,000 sq ft per year) and is prohibited from use in Nassau and Suffolk

Contains LESCO® Poly Plus® Sulfur Coated Urea to provide a uniform growth with extended nitrogen feeding.

ACTIVE INGREDIENT:

Dillriopyr. S.S'-dimethyl 2-(diffuoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

Read the entire label before using this product. Use only according to label instructions. NOTICE: Before using this product, read the Use Precautions, Warranty Statements. Directions for Use, and the Storage and Disposal Instructions. If the Warranty statements are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

# KEEP OUT OF REACH OF CHILDREN CAUTION

	FIRST AND
If swallowed	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
ffin eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue finsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
tf on skin orclotting	
If inhaled	<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! CAUSES EYE IRRITATION. HARNIFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause altergic reaction in some individuals.

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

#### GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass, other annual grasses. and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a Common Name Scientific Name Benigrass, creeping Agrostis palustris Poa prefensis Bluegrass, Kentucky Fescue, the Festuca rubra Festuca arundinacea Fescue, tall Lolium perenne Ryegrass, perennial

p.2

Warm-Season Grasses

Common Name Scientific Name Paspatum notatum Rahiaorass Cyrrodon dactylon Bermudaorass Ruchine decivioldes Euffalograss\*\*⁴ Carpetorass Axonoous affinis Eremochloa ophiuroides Centipedegrass Pennisetum clandestinum Kikuyuprass Stenotaphrum secundalum St. Augustinegrass Zovsia, isponica Zoysiagrass

DO NOT apply this product to Colonial Bentgrass (Agrostis fenuis) varieties.

"Use of this product on certain varieties of Creeping Bentgrass, such as "Cohansey". "Carmen", "Seaside", and Washington may result in undestrable turgrass injury. Not all varieties of Creeping Bentgrass have

"Use of this product on certain varieties of Fine Fescue may result in undesirable turf injury. The following Fine Fescus varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty, 'Bigart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Nelo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Penntawn', 'Tamara', 'Tatjana', 'Waldon', and 'Waldona'. Not all varieties of Fine Fescus have been

\*\*\*DO NOT use this product on seeding Buffalograss in the spring of the first year of establishment unlit the turigrass is fully green and has established new roots.

#### **DIMENSION\* TURFGRASS REGIONS**



Reseeding, Overseeding, or Sprigging

Reseeding, overseeding or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turigrasses. When researing or overseeding, proper cultural practice such as soil cultivation, inigation and fertilization should be followed.

However, if overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application.

For best results, use mechanical or power seeding equipment (stit seeders) designed to give good seed to soil contact.

#### **USE PRECAUTIONS**

- Not for use on Golf Course Putting Greens.
- · The early post-emergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first filler. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product DO NOT apply this product until the grass has recovered from these cultural practices.
- Apply this product directly to established lawns or ornamental turfs only.
- DO NOT apply to flowers, vegetables, shrubs, or trees.
- DO NOT use clippings from treated turf for mulching around vegetables or fruit
- DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled. # 7.4 15 100.004 - 80

# **DIMENSION\* 0.15% Plus Fertilizer**

APPLICATION DIRECTIONS - Control of Crabgrass
Pre-emergence and Early Post-emergence Control

This product provides "pre-emergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turts. It can also provide "early post-emergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turts. Post-emergence crabgrass control will be obtained only when this product is applied prior to the filtering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turt. So the practical benefit of this product's additional, early post-emergence activity is that (compared to strictly pre-emergence crabgrass products). Dimension controls crabgrass prior to and up to 4 weeks after germination.

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

DO NOT apply more than 7.65 to of this product per 1,000 sq ft per application, and no more than 22.95 to of this product per 1,000 sq ft per year.

Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turigrass moving height, and whether the use is considered to be pre-emergence or early post-emergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turigrass competitiveness, and the user's location within a region.

<u>Use Program #1</u> for pre-emergence control at siles where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of pre-emergence crabgrass control. This program should not be used for early post-emergence crabgrass control.

Use Program #2 for pre-emergence control at sites where (a) turigrass is cut low (e.g., gotf fairways), arei (b) tealgrass mainterrance or weed control has not been conducted during the previous year. This program provides 4-6 mornins of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turigrass is cut high (e.g., homeowner lawns).

<u>Use Program #3</u> for pre-emergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut low (e.g., gulf fairways).

Subsequent, sequential pre- and/or post-emergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Table 1: Recommended Single Application Use Rates\*

PROGRAM	#1	#2	#3 Pre-emergence (low-cut turf)	
USE	Pre-emergence	Pre-emergence		
(turforass out)	(high-cut turf)	(low-cut burl)		
(		Post-emergence	Post-emergence	
REGION		(high-cut turt)	(low-cut turf)	
North	1.91 lb/1,000 sq fl <sup>‡</sup>	2.75 Ed/1,000 sq 11 <sup>‡</sup>	3.82 lb/1,000 sq ft	
Transition	2.75 lb/1,000 sq ft <sup>‡</sup>	3.82 th/1,000 sq ft	5.66.b/1,000 sq ft	
South	3.82 lb/1,000 sq ft	5,56 lb/1,000 sq ft	7.65 lb/1,000 sq.ft	
Coastal South	5.66 lb/1,000 sq ft	7.65 fb/1.000 sq ft	7.65 lb/1,000 sq ft**	

Note: 3.82 lb per 1,000 sq ft is equal to 0.25 lb active ingredient per acre.

\*Particle distribution at this rate may not provide adequate control or suppression.

\*DO NOT apply more than 7.65 to of this product per 1,000 sq ft per application, and no more than 22.95 to of this product per 1,000 sq ft per year. Post-emergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below. <u>See</u> man of the United States.

North: all areas not designated below.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH,

& coestal areas of CT, NY, & RI.

South: Ai., AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.
Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

"May require spit or sequential applications for full control.

Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

**Tips for Improved Control** 

For best results, apply this product within a few days after moving and delay moving again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

goosegrass kikuyugrass pursiane (common) speedwell (com) spurge (prostrate & spotted)

\*Also controlled at the "Program 2" rates.

**Fall Applications** 

Used as directed for take summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

bluegrass (annual) geranium (Carolina)
tititercress persiev piert
chickweed pineappleweed
henbit shepherd's purse

SUGGESTED SPREADER SETTINGS\*

	(Recommended Rates in larger 1,000 sq ft)					
Spreader	1.91 15	2.75 to	3.62 %	5.65 lb	7.65 to	
LESCO Calibration Gauge	12	13	15	17	20	
SCOTTS* R8A	1	1/2	1	КЖ	M	
Declare or Several	3%	3%	4	41/2	4%	
LESCO Pendukm	- 18	20	24	32	38	

"IMPORTANT: These settings are only approximate: Age, condition of spreader, and operator speed can cause wide variation. Be sure to cathonic your spreader with each application.

#### **APPLICATION EQUIPMENT AND INSTRUCTIONS**

Apply this product with drop or rotary spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturier's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning property and applying uniform distribution of granules.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secured storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: Nonrefillable container. Do not resse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPILL: In case of spill, sweep up material and dispose of material according to "Product Disposal" directions listed above.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

MOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be returned.

The Directions for Use of this product must be informed carefully and completely. It is impossible to eliminate all risks inhanently associated with the use of this product. To the extent consistent with applicable law. Buyer analon User assume all risks of ineffectiveness or other unintended consequences or demands that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the fine period in which the product is applied, the presence of other makerials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc. To the extent consistent with applicable law, all such risks shall be assumed by Buyer analon User, and Buyer analon User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product conting to labellinshustors, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, Inc. MAYES NO WARRANTES OF NERCHANTABLITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR MAY LED WARRANTY EXCEPT AS STATED ABOVE.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INURIES OR DAMACES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, REGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR F NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, INC., BE LABLE FOR ANY INCIDENTAL, CONSEQUENTAL OR SPECIAL DAMAGES.

LESCO, Inc. offers this product, and Buyer another User accepts 4, subject to the tonegoing Conditions of Sale and Limitation of Watmarky and Lishbity, which may not be modified except by written agreement signed by a dust authorized representative of LESCO, Inc.